

Case Report

Open Access

Complete heart block and severe aortic stenosis in a patient with rheumatoid arthritis: a case report

Ioannis Moysakis, Nikolaos Lionakis*, Ioannis Vlahodimitris and Vassilios Votteas

Address: Laiko General Hospital of Athens, Department of Cardiology, 17 Agiou Thomas St, GR-15727 Goudi-Athens, Greece

Email: Ioannis Moysakis - moysak@yahoo.gr; Nikolaos Lionakis* - nik_lion@yahoo.com; Ioannis Vlahodimitris - dr_eam_74@hotmail.com; Vassilios Votteas - focus_dd@hotmail.com

* Corresponding author

Published: 5 February 2009

Received: 7 November 2008

Cases Journal 2009, 2:126 doi:10.1186/1757-1626-2-126

Accepted: 5 February 2009

This article is available from: <http://www.casesjournal.com/content/2/1/126>

© 2009 Moysakis et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: A 77-year-old male patient with a history of rheumatoid arthritis was admitted to our hospital for investigation of syncope and dyspnea on exertion class II according to NYHA class association.

Case presentation: The electrocardiogram revealed complete heart block whereas the echocardiogram showed severe aortic valve stenosis with a peak gradient = 80 mmHg. A permanent pacemaker was implanted in addition to aortic valve replacement. The coexistence of complete heart block and severe aortic stenosis with rheumatoid arthritis are presented.

Conclusion: Further studies are necessary to assess whether a true association of the above conditions exist.

Background

Rheumatoid arthritis is a chronic autoimmune multisystem disease which can affect the pericardium, myocardium, and endocardium. Heart involvement is a frequent cause of death [1-3].

This report presents a case with severe aortic valve stenosis combined with complete heart block developing in a patient with rheumatoid arthritis. It would be interesting to see whether a linkage between the above cardiac lesions with rheumatoid arthritis can be demonstrated.

Case report

A 77-year-old male with a known history of rheumatoid arthritis was admitted to our hospital due to syncopal epi-

sodes and dyspnea on exertion class II according to (NYHA). Rheumatoid arthritis was diagnosed 39 months ago on the basis of clinical and laboratory findings [4]. The patient was given corticosteroids (7.5 mg prednisone daily).

On physical examination her heart rate was 42 beats per minute and blood pressure 135/80 mmHg. Auscultation revealed a 3/6 systolic murmur at the right second intercostal space which radiated to the carotids and significant diminished the second heart sound. Laboratory findings were as follows: hematocrit 34.9%, white cell count 8.800/mm³, platelet count 254000/mm³, erythrocyte sedimentation rate 57 mm/1st h, CRP 10.5 mg/L urea 53 mg/dl, creatinine 1.1 mg/dl, SGOT 17, SGPT 24 U/L. Ra-test

was negative whereas disease activity index and health assessment questionnaire score were 4.9 and 1.4 respectively.

A resting electrocardiogram showed complete heart block (figure 1). Transthoracic echocardiography demonstrated severe left ventricular hypertrophy. The aortic valve was tricuspid with calcific and thickened aortic leaflets causing decreased opening. A transvalvular peak gradient of 80 mmHg was measured by continuous Doppler whereas a mild degree aortic regurgitation was revealed. Due to the fact that the ejection fraction was 45% the aortic valve orifice was measured which was 0.7 cm². The patient sustained a coronary angiogram which was negative for coronary artery disease. Therefore, a permanent DDD pacemaker was implanted in addition to aortic valve replacement.

Discussion

Rheumatoid arthritis is a common chronic autoimmune disease involving many organ systems, primarily the joints; frequently it is accompanied by cardiac lesions including pericardium, myocardium and endocardium, comprising coronary arteries, valvular tissue and conduction system. Heart involvement in RA is the leading cause of death [1-3].

Conduction disturbances are well recognized in the rheumatoid arthritis population. Primary infiltration by mononuclear cells or rheumatoid nodules could cause conduction disturbances at the sino-atrial or atrioventricular nodes or His-Purkinje system [5,6]. Other potential mechanisms are vasculitis of the arterial supply to conductive tissue, haemorrhage into a rheumatoid nodule or extension of an inflammatory lesion from the aortic or mitral valve [6]. Rarely, these lesions may be due to amy-

loid deposition [6]. Ahern et al have described complete heart block in 8 patients with rheumatoid arthritis and reviewed 20 similar cases previously reported.[7] However complete atrioventricular block is rare in RA and does not respond to anti-inflammatory and immunosuppressive therapy [6].

Valvular heart disease associated with rheumatoid arthritis has not been well characterized and its clinical predictors are undefined [8]. The incidence of rheumatoid valvular disease is among 2.5%–30% of necropsy patients with the above disease [9]. Valve nodules and valve thickening diffuse or localized by unspecific inflammatory changes were found in aortic and mitral valves [8,9]. No correlation was also found between valvular disease and duration, activity, severity, pattern of onset and course, extra-articular disease, serology, or therapy of RA [8].

Our case is interesting since there is no clear explanation for the severe valvular lesion combined with complete heart block in a patient with rheumatoid arthritis. In spite of the fact that the systemic disease was mild in severity complete heart block and severe aortic stenosis occurred. It should be noted that the patient had no history of arterial hypertension or renal involvement, conditions associated with aortic valve calcification. Moreover our case was not received anti-TNF and specifically infliximab therapy which are recognized causing atrioventricular block [10]. The existent severe aortic valve stenosis possibly could be associated with the atrioventricular block.

To our knowledge there are no reports in the medical literature of complete heart block and severe aortic valve stenosis in patients with rheumatoid arthritis. The association of the above cardiac lesions with rheumatoid arthritis may, however, be a random occurrence.




Figure 1
The ECG of the patient showing the complete atrioventricular heart block.

Consent

Written informed consent was obtained from the patient for publication of this case report and any accompanying images. A copy of the written consent is available for review by the Editor-in-Chief of this journal.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

IM conceived the case report. NL and IV were involved in the case management, and drafted the manuscript. IM, NL and VV reviewed the manuscript and made the final corrections before submission. All the authors have read and approved the final manuscript.

References

1. Lebovitz WB: **The heart in rheumatoid arthritis (rheumatoid disease). A clinical and pathological study of sixty-two cases.** *Ann Intern Med* 1963, **58**:102-123.
2. Hernandez-Lopez E, Chahine RA, Anastasiades P, Raizner AE, Lidsky MD: **Echocardiographic study of the cardiac involvement in rheumatoid arthritis.** *Chest* 1977, **72**:52-55.
3. Guedess C, Bianchi-Fior P, Cormier B, Barthelemy B, Rat AC, Boissier MC: **Cardiac manifestations of rheumatoid arthritis: A case-control transesophageal echocardiography study in 30 patients.** *Arthritis Rheum* 2001, **45**:129-35.
4. Arnett FC, Edworthy SM, Block DA, McShane DJ, Fries JF, Cooper NS, et al.: **The American Rheumatism Association 1987 revised criteria for the classification of Rheumatoid Arthritis.** *Arthritis Rheum* 1988, **31**:315-24.
5. Pizzarello RA, Goldberg J: **The heart in rheumatoid arthritis.** In *Rheumatoid arthritis: aetiology, diagnosis and management* Edited by: Utsinger PD, Zvaifler NJ, Ehrlich GE. Philadelphia: J B Lippincott; 1985:431-40.
6. Seferovic PM, Ristic AD, Maksimovic R, Simeunovic DS, Ristic GG, Radovanovic G, et al.: **Cardiac arrhythmias and conduction disturbances in autoimmune rheumatic diseases.** *Rheumatology (Oxford)* 2006, **45**(Supplement 4):39-42.
7. Ahern M, Lever JV, Cosh J: **Complete heart block in rheumatoid arthritis.** *Ann Rheum Dis* 1983, **42**:389-97.
8. Roldan CA, DeLong C, Qualls CR, Crawford MH: **Characterization of valvular heart disease in rheumatoid arthritis by transesophageal echocardiography and clinical correlates.** *Am J Cardiol* 2007, **100**:496-502.
9. Asin Ungria J, Pina Salvador JM, Beltran Audera J, Otero Escalante W, Cia Gomez P: **Double heart prosthesis in a patient with rheumatoid arthritis.** *An Med Interna* 1989, **6**:33-5.
10. Sote Y, Green S, Maddison P: **Complete heart block after infliximab therapy.** *Rheumatology (Oxford)* 2007, **47**:227-28.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

