

**Oral Squamous Cell Carcinoma with an unusual clinical manifestation: a case report**Farnaz Falaki<sup>1\*</sup>, Zahra Delavarian<sup>2</sup>, Atessa Pakfetrat<sup>1</sup>, Nooshin Mohtasham<sup>3</sup> and Shiva Shirazian<sup>4</sup>

Addresses: <sup>1</sup>Department of Oral Medicine, Faculty of Dentistry and Dental research center, Vakilabad Blvd, Mashhad, postal code:91735, Iran, <sup>2</sup>Department of Oral Medicine, Faculty of Dentistry and Dental research center, Mashhad University of Medical Sciences, postal code:91735, Iran, <sup>3</sup>Department of Oral and Maxillofacial Pathology, Faculty of Dentistry and Dental research center, Mashhad University of Medical Sciences, Mashhad, postal code:91735, Iran and <sup>4</sup>Specialist in Oral Medicine, No.32-Shariati street-Tehran-Iran

Email: FF\* - falakif@mums.ac.ir or farnazfalaki@yahoo.com; ZD - Delavarianz@mums.ac.ir; AP - pakfetrata@mums.ac.ir; NM - mohtashamn@mums.ac.ir; SS - shiraziash@gmail.com

\* Corresponding author

Published: 20 April 2009

Received: 22 December 2008

Cases Journal 2009, 2:6608 doi: 10.1186/1757-1626-2-6608

Accepted: 9 February 2009

This article is available from: <http://casesjournal.com/casesjournal/article/view/2/4/6608>

© 2009 Falaki et al; licensee Cases Network Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

**Abstract**

**Introduction:** Squamous cell carcinoma is the most common malignant tumor of the oral cavity and one of the 10th most common causes of death. It arises from dysplastic oral squamous epithelium. By considering the pathogenesis of squamous cell carcinoma, the smooth and intact surface for this lesion is not usual.

**Case presentation:** A painful nodular lesion with smooth surface on the left buccal mucosa of a 75-year-old female patient was observed. She noticed it 2 weeks ago. Histopathological examination revealed oral squamous cell carcinoma.

**Conclusion:** In this paper, we report an unusual clinical presentation of oral squamous cell carcinoma in buccal mucosa which is very rare.

**Introduction**

Approximately 94% of all oral malignancies are Squamous cell carcinoma (SCC). The annual incidence and mortality rates vary between different races, genders, and age groups. In the United States this is 7.7 per 100,000 [1].

Like other carcinomas, the risk of intra oral carcinoma increases with increasing age especially for males [1-4]. Persons with oral SCC almost have been aware of an alteration in that site for 4-8 months before seeking

professional help. There is minimal pain during the early growth phase and this may explain the delay in seeking professional care [1].

If the health care professional does not have a high index of suspicion, additional several weeks or months may elapse before a biopsy is performed [1].

Oral SCC has various clinical presentations such as exophytic, endophytic, leukoplakic and erythroplakic, which all of them show visible changes in the surface.

In the present paper, we report a case of exophytic oral SCC with a smooth surface which is an unusual presentation [1,5].

### Case presentation

A 75-year-old Caucasian female of Iranian nationality was admitted to the Department of Oral Medicine in Mashhad Dental Faculty with chief complaint of a painful mass in the left buccal mucosa, which was first noticed by the patient 2 weeks earlier and gradually increased in size.

Intra-oral examination revealed a painful normal-colored nodular lesion with smooth surface in the left buccal mucosa with small yellow papules at the surface with approximate size of 2.5 × 1.5 cm and firm in consistency (Figure 1).

The patient did not feel any sensory changes in the affected area. Physical examination revealed no lymphadenopathy in submandibular or other neck triangles.

The patient was diabetic (type II). She did not have any risk factor for SCC (smoking or alcohol consumption) and have had 4 pregnancies.

By considering the characteristics of the lesion and our physical examination findings, our differential diagnosis

were minor salivary gland tumors and other tumors of mesenchymal origin.

Incisional biopsy under local anesthesia was performed and the specimen was submitted for histopathological examination, which revealed a malignant neoplastic proliferation of stratified squamous epithelial cells as sheets or islands of cells, invading to the connective tissue (Figure 2).

The definite diagnosis was squamous cell carcinoma (Grade I). The patient was referred to an oncology department where surgery by using intra oral approach was performed and diagnosis of SCC was reconfirmed, but she died 3 days after surgery because of poor management of diabetes.

### Discussion


This is an unusual case of oral SCC in a 75-year-old female patient with a nodular lesion having a smooth and intact surface.

By considering the pathogenesis of SCC all presentations are associated with changes in the surface as expected for epithelial lesions and all exophytic SCCs have a rough surface and irregular shape.

On very rare occasion, squamous cell carcinoma may commence at a small location on the surface, burrow and undermine the subepithelial tissue in such a manner that the lesion appears mostly as a smooth surfaced exophytic




**Figure 1.**  
clinical view of the exophytic lesion with smooth surface in the buccal region. Small yellow papules are visible at the surface.




**Figure 2.**  
Invasion of tumor cells to connective tissue, Magnification 40 X, Hematoxylin & Eosin staining.






**Figure 3.**  
Pleomorphism, Hyperchromatism, and keratin formation in tumor, Magnification 100 X, Hematoxylin & Eosin staining.



**Figure 4.**  
Pleomorphism and invasion of tumor cells, Magnification 400 X, Hematoxylin & Eosin staining.



**Figure 5.**  
Smooth and intact epithelium of the lesion, tumor cells in connective tissue, Magnification 40 X, Hematoxylin & Eosin staining.

lesion, which makes a diagnostic challenge [6]. To the best of our knowledge it is the only reported case of oral SCC with smooth and intact surface reported in the literature.

The purpose of this article is to emphasize that even in smooth-surfaced rapid-growing oral lesions, SCC should be considered in differential diagnosis and this needs a careful examination and management by both medical and dental practitioners.

#### Abbreviation

SCC: Squamous cell carcinoma.

#### Consent

Written informed consent was obtained from the patient for publication of this case report and accompanying images. A copy of written consent is available for review by the Editor-in-Chief of this journal.

#### Competing interests

The authors declare that they have no competing interest.

#### Authors' contributions

FF, ZD and SS analyzed and interpreted the patient's data and performed the clinical examination, NM performed the histological examination of the lesion, AP assisted in clinical study and writing the manuscript. All authors read and approved the final manuscript.

## Acknowledgement

The authors are supported by dental research center of Mashhad Dental Faculty and the vice chancellor for research of Mashhad university of Medical sciences (MUMS).

## References

1. Neville BW, Damm DD, Allen CM, Bouquot JE: **Oral & Maxillofacial Pathology**. 2nd ed. Philadelphia: VVB Saunders; 2002: 356-366.
2. Greenberg MS, Glick M: **Burket's Oral Medicine**. 10th ed. BC Decker Inc, 2008; 153-160.
3. Seoane J, Warnakulasuriya S, Varela-Centelles P, Esparza G, Dios PD: **Oral cancer: experiences and diagnostic abilities elicited by dentists in North-Western Spain**. *Oral Disease* 2006, **12**(5):482-492.
4. Lawoyin JO, Lawoyin Do, Fasola Ao, Kolude B: **Intra-oral squamous cell carcinoma in Nigerians under 40 years of age: a clinicopathological review of eight cases**. *Afr J Med Med Sci*. Mar 2005, **34**(1):99-102.
5. Regezi JA, Sciubba JJ, Gordan R CK: **Oral Pathology**. 4th ed. Saunders; 2008:48-54.
6. Wood NK, Goaz PW: **Differential Diagnosis of Oral and Maxillofacial Lesions**. Fifth ed. Mosby; 1997:137.

## Do you have a case to share?

Submit your case report today

- Rapid peer review
- Fast publication
- PubMed indexing
- Inclusion in Cases Database

**Any patient, any case, can teach us something**



**CASES  
NETWORK**

[www.casesnetwork.com](http://www.casesnetwork.com)