



Case report

***Mycobacterium avium-intracellulare* infection presenting as a testicular mass in an immunocompromised patient: a case report**

Ruth A Hartley

Address: Department of Haematology, East Surrey Hospital, Canada Avenue, Redhill, Surrey, RH1 5RH, England

Email: ruthhartley@doctors.net.uk

Received: 2009 Accepted: 2009 Published: 19 August 2009

Cases Journal 2009, **2**:8975 doi: 10.4076/1757-1626-2-8975

This article is available from: <http://casesjournal.com/casesjournal/article/view/8975>

© 2009 Hartley; licensee Cases Network Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Isolated mycobacterial infections of the genital tract are extremely rare. Here we present a 63-year-old gentleman who developed an isolated non-tuberculous mycobacterial mass in the inferior pole of his right testis, secondary to immunocompromise related to multiple autoimmune and haematological conditions and their treatments.

Introduction

Compromised immunity, associated with autoimmune and haematological conditions and their treatments, causes an estimated increase of 3.69 infections per patient, with a mortality of 21% [1]. Autoimmune conditions are also linked to the development of lymphoma, with a 3.16 relative risk of lymphoma developing in patients with systemic lupus erythematosus (SLE) [2]. The crossover between these two conditions can be profound [3], and the diagnostic intricacies can become even more complex given the prevalence of atypical infections which can mimic relapse of either condition.

Of the atypical infections, tuberculous and non-tuberculous mycobacteria (NTM) are common pathogens, most frequently in the chest, but exceptionally affecting the genitourinary tract [4]. Here we present a patient with an isolated NTM infection of his right testis due to immunocompromise from SLE, recent lymphoma, and immunosuppressive drugs.

Case presentation

Our patient was a 63-year-old retired gentleman, weighing 64 kilograms and 1.8 meters tall. He had no significant family history and drank only occasionally. At presentation he was trying to give up a 40 pack-year smoking habit. His medications were predominately analgesics but also a bisphosphonate and prednisolone 20 mg.

He had a history of SLE, haemolytic anaemia, recurrent idiopathic thrombocytopenia purpura and recent chemotherapy (6 months previously) for diffuse large B-cell lymphoma.

He presented to clinic with a tender, diffusely enlarged right testicle, and was also noted to be suffering from night sweats, a 7 kg weight loss, and splenomegaly (previously noted prior to treatment for lymphoma).

Tumour markers were normal, but an ultrasound scan of his right testicle showed a non-homogenous abnormality

inferiorly (Figure 1), in keeping with a recurrence of lymphoma or a primary malignancy [5].

Given his history and symptoms, he was urgently referred to the urologists who decided to proceed with a right orchidectomy.

Histology of the right testis showed a granulomatous orchiditis. A Ziehl-Neelson (ZN) stain showed a large number of acid fast bacilli (AFB) which, given the morphology and absence of caseation, was most consistent with a NTM infection. Unfortunately, no sample had been sent for culture, but given the histological findings and clinical presentation, this was considered most likely to be *Mycobacterium avium-intracellulare* (MAI) infection.

He recovered well post operatively, and, as a CT scan showed a right apical lung lesion, he was referred to the local respiratory physician for further investigation and management.

Bronchial washings of the right upper zone and left lingula were positive for AFB. He was commenced on conventional

quadruple anti-tuberculous therapy pending formal identification of the AFB. The organism was subsequently confirmed as MAI, and therapy was switched according to the British Thoracic Society guidelines [6].

Discussion

NTM infections in immunocompromised patients [7] are often associated with disseminated disease, with the chest typically as the primary focus of infection [8]. The genitourinary system can become involved, as with tuberculous disease, but involvement of the testes without early epididymal changes, as occurred in this case, was previously considered as almost pathognomonic of a neoplastic rather than infectious process [1,9].

Conclusion

Mycobacterial infections in immunocompromised patients must be excluded due to the importance in prognosis and treatment (early treatment of MAI results in remission rates of 67% even in the immunocompromised) [10]. In this gentleman, his extreme immunocompromised state made him the ideal candidate for opportunistic infections, resulting in the situation where unusual presentation of an infection mimicked a recurrence of lymphoma.

Consent

Written informed consent was obtained from the patient for publication of this case report and accompanying images. A copy of the written consent is available for review by the Editor-in-Chief of this journal.

Competing interests

The author declares that she has no competing interests.

Acknowledgements

Dr Ann Nandi (Consultant Haematologist, East Surrey Hospital), Dr Warren Shattles (Consultant Rheumatologist, East Surrey Hospital), Dr Bruce Stewart & Dr Karen Knox (Consultant Microbiologists, East Surrey Hospital), Dr Ali Alhakim (Consultant Histopathologist, East Surrey Hospital), Dr Iain Wilkinson (ST2 Haematology, East Surrey Hospital).

References

1. Cohen J, Pinching A, Rees A, Peters D: **A study of the infective complications of 75 patients with immunologically-mediated disease.** *Q J Medicine* 1982, **51**:1-15.
2. Bernatsky S, Ramsey-Goldman R, Isenberg D, Rahman A, Dooley M, Sibley J *et al.*: **Hodgkin's lymphoma in systemic lupus erythematosus.** *Rheumatology* 2007, **46**:830-832.
3. Bernatsky S, Ramsey-Goldman R, Lachance S, Pineau C, Clarke A: **Lymphoma in a patient with systemic lupus erythematosus.** *Nat Clin Pract Rheumatol* 2006, **2**:570-574.
4. Drudi FM, Laghi A, Iannicelli E, Di Nardo R, Occhiato R, Poggi R *et al.*: **Tubercular epididymitis and orchitis: US patterns.** *Eur Radiol* 1997, **7**:1076-1078.
5. Muttarak M, Peh W: **Tuberculous Epididymo-orchitis.** *Radiology* 2006, **238**:748-751.




Figure 1. Ultrasound scan of the right testicle showing the non-homogenous abnormality inferiorly (white arrow).

6. **Subcommittee of the Joint Tuberculosis Committee of the British Thoracic Society: Management of opportunist mycobacterial infections: Joint Tuberculosis Committee Guidelines.** *Thorax* 2000, **55**:210-218.
7. Field S, Cowie R: **Lung disease due to the more common nontuberculous Mycobacteria.** *Chest* 2006, **129**:1653-1672.
8. Corti M, Palmero D: **Mycobacterium avium complex infection in HIV/AIDS patients.** *Expert Review of Anti-infective Therapy* 2008, **6**:351-363.
9. Senzaki H, Watanabe H, Ishiguro Y: **A case of very rare tuberculosis of the testis [abstract].** *Nippon Hinyokika Gakkai Zasshi* 2001, **92**:534-537 [Japanese].
10. Field S, Cowie R: **Treatment of mycobacterium avium-intracellulare complex lung disease with a macrolide, ethambutol, and clofazimine.** *Chest* 2003, **124**:1482-1486.

Do you have a case to share?

Submit your case report today

- Rapid peer review
- Fast publication
- PubMed indexing
- Inclusion in Cases Database

Any patient, any case, can teach us something



**CASES
NETWORK**

www.casesnetwork.com