

## Case report

**Transient lung herniation through a thoracic cage defect:  
a case report**

Kostas Psathakis\*, Charalampos Mermigkis and Kostas Tsintiris

Address: Department of Pneumology, Army General Hospital of Athens, Greece

Email: KP\* - kpsazakis@hol.gr; CM - mermigis@hotmail.com; KT - kostastsintiris@yahoo.gr

\* Corresponding author

Received: 18 September 2008 Accepted: 4 April 2009 Published: 12 June 2009

*Cases Journal* 2009, **2**:7524 doi: 10.4076/1757-1626-2-7524This article is available from: <http://casesjournal.com/casesjournal/article/view/7524>

© 2009 Psathakis et al; licensee Cases Network Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.**Abstract**

We report a benign condition of transient lung herniation through a congenital structural defect of the thoracic cage, in a young, otherwise healthy, asymptomatic individual. A brief review of the existing literature on this rare entity is also presented.

**Case presentation**

A 23-year-old healthy Greek male, came to the outpatient clinic of our hospital for a routine examination for employment. During the examination, he mentioned that his right chest wall protruded whenever he performed a Valsalva's maneuver (Figure 1). He said that it happened since he was a child and he had been informed that it was a benign condition. The rest of the clinical examination, pulse oximetry as well as spirometry were normal. His chest radiography revealed that the 8<sup>th</sup> and 9<sup>th</sup> ribs on the right departed from each other leaving a gap (Figure 2). This configuration could explain the transient lung herniation that happened whenever the intrathoracic pressure was increased (e.g. during cough, Valsalva's maneuver etc).

As the patient was asymptomatic, no further intervention was considered.

**Discussion**

Herniation of the lung is defined as a protrusion of the lung beyond the normal confines of the thoracic cavity through

an abnormal opening of the chest wall [1]. In reference to the underlying mechanism, a sudden growth of intrathoracic pressure is in the background, which can be caused by coughing, emesis, lifting heavy weights; usually combined with attenuation of the thoracic wall [2].


The first case of lung herniation was recorded by Roland in 1499. Since then about 300 patients have been reported, most of them as single case reports. In the majority of cases, the lung herniated through the intercostal space as a result of trauma or after thoracic operation; most of the other hernias were congenital [1,3].

The most widely accepted classification of lung hernia is that of Morel-Lavallee, based on both the anatomic location and etiology. According to their anatomic location pulmonary hernias are classified as diaphragmatic, thoracic (intercostal) and cervical. According to their etiology they are classified as congenital or acquired. Acquired hernias are further classified as spontaneous, traumatic and pathological (secondary to inflammatory or neoplastic processes) [1,4].



**Figure 1.** A Valsalva's maneuver causes herniation of the right lung through a "gap" of the thoracic cage.

A spontaneous lung herniation is more frequent in men and in patients with chronic obstructive pulmonary disease, presumably as a result of chronic coughing and hyperinflation, perhaps combined with long-term steroid



**Figure 2.** A chest radiography reveals that the anterior portion of the 8<sup>th</sup> rib comes closer to the 7<sup>th</sup> rib leaving a gap between the 8<sup>th</sup> and 9<sup>th</sup> ribs (white arrows). Note that the 5<sup>th</sup> right rib has an abnormal configuration as well (black arrow).

administration. The anterior thorax, between the 8<sup>th</sup> and 9<sup>th</sup> ribs is the site of predilection for this type of hernias, presumably because of the diminished muscular support at this region [4]. It might be interesting to observe that our patient had a congenital gap of the thoracic cage at this particular intercostal space. The gap combined with the low resistance of the chest wall at this area give a better explanation of the underlying mechanism of the transient lung herniation in this patient.

Postoperative intercostal hernia can develop through disjoining of the intercostal space on the area of thoracotomy or video thoracoscopy [2].

Most lung hernias are asymptomatic. It is likely that a proportion of cases goes unnoticed or gives rise to only minimal symptoms that do not require medical consultation, and therefore the true incidence of the condition is difficult to estimate. This could explain why most pulmonary hernias reported in the literature are not spontaneous but are related to various traumatic conditions [1]. Symptomatic patients usually present with a bulging, crepitant mass protruding through the chest wall either with or without pain. The most important clue is exacerbation of the hernia on Valsalva maneuver, coughing, or straining, with resolution on inspiration or quiet breathing [5].

Incarceration and strangulation, pain, hemoptysis or recurrent infection are the possible complications of the herniation but are rarely observed. In asymptomatic cases conservative management may be sufficient with compressive pads and corsets, and treatment of the underlying condition causing increased intrathoracic pressure. Surgery offers a definitive treatment and is recommended in case of symptoms, interference with everyday activities or for cosmetic reasons [1,4]. It is done by the contraction of the intercostal space with percostal stitches and the reconstruction of the musculature or the implantation of plastic surgical mesh, to cover the gap [2].

In the described patient, the volume of the herniated lung was probably not large enough to affect his every day activities or spirometry. Moreover, since he was a healthy individual with a normal musculature, the lung herniated transiently only when the intrathoracic pressure exceeded the resistant of the muscles that covered the congenital gap of the chest wall. Perhaps for these reasons he had no symptoms since his childhood. Although some authors recommend surgical repair of the defect even in asymptomatic patients [4], we suggest that asymptomatic lung hernias can be left untreated if they do not cause any problems. Obviously, these patients should be followed up and any intervention is justified when symptoms occur [3,5].

### Consent

Written informed consent was obtained from the patient for publication of this case report and accompanying images. A copy of the written consent is available for review by the Editor-in-Chief of this journal.

### Competing interests

The authors declare that they have no competing interests.

### Authors' contributions

KP examined the patient and wrote the manuscript. CM examined the patient and processed the photos. KT had the overall responsibility of the clinic and the final approval of the manuscript.

### References

1. Tack D, Wattiez A, Schtickzelle JC, Delcour C: **Spontaneous lung herniation after a single cough.** *Eur Radiol* 2000, **10**:500-502.
2. Szentkereszty Z, Boros M, Sápy P, Kiss SS: **Surgical treatment of intercostal hernia with implantation of polypropylene mesh.** *Hernia* 2006, **10**:354-356.
3. Weissberg D, Refaely Y: **Hernia of the lung.** *Ann Thorac Surg* 2002, **74**:1963-1966.
4. Sulaiman A, Cottin V, Pereira de Souza Neto E, Orsini A, Cordier JF, Gamondes JP, Tronc F: **Cough induced intercostal lung herniation requiring surgery: report of a case.** *Surg Today* 2006, **36**:978-980.
5. Ross RT, Burnett CM: **Atraumatic lung hernia.** *Ann Thorac Surg* 1999, **67**:1496-1497.

### Do you have a case to share?

Submit your case report today

- Rapid peer review
- Fast publication
- PubMed indexing
- Inclusion in Cases Database

**Any patient, any case, can teach us something**



**CASES  
NETWORK**

[www.casesnetwork.com](http://www.casesnetwork.com)