# Cases Journal



Case Report Open Access

# A rare presentation of appendicitis as groin swelling: a case report Maheswaran Pitchaimuthu\*1,2 and Stephen Dace1,3

Address: <sup>1</sup>Department of General Surgery, Altnagelvin Area Hospital, Londonderry, Northern Ireland, UK, <sup>2</sup>SpR in HPB Surgery, Imperial College Healthcare NHS Trust, Hammersmith Hospital, London, W12 0HS, UK and <sup>3</sup>Consultant Surgeon, Western Health and Social Care Trust, Altnagelvin Hospital, Londonderry, BT47 6SB, UK

Email: Maheswaran Pitchaimuthu\* - drmpitchaimuthu@yahoo.com; Stephen Dace - Stephen.Dace@westerntrust.hscni.net

\* Corresponding author

Published: 14 January 2009

Cases Journal 2009, 2:53 doi:10.1186/1757-1626-2-53

This article is available from: http://www.casesjournal.com/content/2/1/53

© 2009 Pitchaimuthu and Dace; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<a href="http://creativecommons.org/licenses/by/2.0">http://creativecommons.org/licenses/by/2.0</a>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Received: 12 October 2008 Accepted: 14 January 2009

**Abstract** 

Background: Appendicitis in femoral hernia is a rare condition, which raises diagnostic challenge.

Case Report: A 40-year-old man presented with painful right-sided groin swelling of I-week duration. The area was explored, with presumpted diagnosis of inguinal abscess. At exploration a femoral hernia was found which contained a mildly inflammed appendix. Appendicectomy and hernia repair was done. Post surgical course was uneventful. We present this case with brief summary of literature pertaining to such lesions.

**Discussion:** The rare occurrence of femoral hernia containing appendix may be explained by different degrees of intestinal rotation during development or variation in its attachment to the caecum. Inflammation is due to tight femoral ring. Preoperative diagnosis is difficult. Management options are diverse.

Conclusion: We present this case because of rarity. Early surgery prevents complications.

# Introduction

Inflammatory swellings of the groin are common, and the changes are often attributed to infection. Although this is possible, inflammatory swellings are often secondary to groin hernia. We present an unusual case of groin swelling, outlining its investigations and subsequent management.

# Case report

An otherwise healthy 40-year-old male presented with a one-week history of pain and swelling in his right groin. There was no history of trauma or previous hernia, and his bowel habit was normal. On examination he had a right inguinal swelling ( $6 \times 3$  cm) lateral to the pubic tubercle. There was no evidence of a cough impulse.

An ultrasound of the region was performed (Figure 1), which showed evidence of cellulitis and a fluid collection. The fluid was aspirated; it was blood stained with no evidence of pus. Routine blood tests were normal.

Given the clinical findings, he was taken to the operating theatre for exploration. A standard oblique groin incision was used, and an incarcerated femoral hernia was identified. The back of the inguinal canal and the neck of the sac were opened. Inside the sac, a long, mildly inflamed appendix was found (Figure 2). An appendicectomy was performed and the excess sac excised and transfixed. Given that there was minimal inflammation of the appendix and there was no obvious evidence of infection outside the sac, the back wall of the inguinal canal was



Figure I USS right groin showing fluid collection.

repaired using the Lichtenstein tension-free mesh method, using  $15 \times 7$  cm Vypro Mesh and 2-0 prolene. The patient had an uneventful recovery.

# **Discussion**

The main pathologic conditions manifesting as masses in the groin fall into five major groups: congenital abnormalities, non-congenital hernias, vascular conditions, infectious or inflammatory processes, and neoplasms [1,2]. The hernial sac may contain preperitoneal fat, omentum, colon, or small bowel but reports of femoral hernia containing the vermiform appendix are rare, reported to occur in 0.8% of femoral hernia [3]. Rence Jacques Croissant de Garengeot, an 18<sup>th</sup> century Paris Sur-




Figure 2
Intra operative picture showing appendix in femoral hernia sac.

geon, was the first to describe appendix in femoral hernia [4]. Appendix in left femoral hernia [5], carcinoid tumour of appendix [6] and stomach [7] as femoral hernia contents have also been reported in the past. Factors contributing to this condition include degrees of intestinal malrotation or the presence of an abnormally large caecum which extends into the pelvis [3].

It is very difficult to diagnose the presence of an appendix within a femoral hernial sac, and to date, only one case has been diagnosed prior to surgery. This was identified at CT scan [8].

Tight femoral ring leads to strangulation and appendicitis. Appendicitis in a femoral hernia does not usually lead to abdominal peritonitis, due to narrow hernia sac neck which prevents inflammation of the parietal peritoneum. Clinical signs include local groin swelling, inflammation and spreading cellulitis, but often the patient feels generally well with no systemic features of sepsis, as in this case [3].

If left untreated, the inflammation may resolve or lead to complications including abscess [9], necrotizing fasciitis [10], necrosis of hernial contents [11] and development of bowel obstruction [12] and even death.

Due to the rarity of such cases treatment options remain diverse. Each case should be judged separately, and treatment based on the principles of removing the source of sepsis (either operatively or by aspiration) should be employed [13,14].

# **Conclusion**

We present a case of acute appendicitis complicating an incarcerated femoral hernia. As is often the case, the diagnosis was made at surgery. By following the principles of removing the source of sepsis and repairing the hernial defect, the patient made a safe recovery from a potentially serious condition.

We present this case because of appendix in femoral hernia that too in a young male patient is very rare. Early surgical treatment prevents potential complications.

# Consent

A written informed consent was obtained from the patient for publication of this case report and accompanying images. A copy of the written consent is available for review by the Editor in Chief of this journal.

# **Competing interests**

The authors declare that they have no competing interests.

### **Authors' contributions**

MP was the oncall registrar, who performed the surgery and a primary author for this manuscript. SD was the oncall consultant who was a contributor in writing the manuscript.

#### References

- Shadbolt CL, Heinze SBJ, Dietrich RB: Imaging of Groin Masses: inguinal anatomy and pathologic conditions revisited. Radio-Graphics 2001, 21:s261-71.
- Apostolidis S, Papavramidis TS, Michalopoulos A, Papadopoulos VN, Paramythiotis D, Harlaftis N: Groin Swelling, the Anatomic Way Out of Abdominal Haematomas: a Case Report and Explicative Literature Review. Acta Chir Belg 2008, 108:251-253.
- Nguyen ET, Komenaka IK: Strangulated femoral hernia containing a perforated appendix. Can | Surg 2004, 47(1):68-9.
- Akopian G, Alexander M: De Garengeot hernia: appendicitis within a femoral hernia. Am Surg 2005, 71(6):526-7.
- Scepi M, Richer JP, Muller J: Appendix in a left crural herniated position: apropos of a case. Explanation by human ontogenesis. | Chir (Paris) 1993, 130(11):479-82.
- lvicic J, Zaloudik J: Carcinoid of the appendix in incarcerated femoral hernia. Rozhl Chir 1999, 78(7):359-61.
- Isaacs LE, Felsenstein CH: Acute appendicitis in a femoral hernia: an unusual presentation of a groin mass. J Emerg Med 2002, 23(1):15-8.
- Lane MJ, Liu DM, Huynh MD, Jeffrey RB Jr, Mindelzum RE, Kats DS: Suspected acute appendicitis: non enhanced helical CT in 300 consecutive patients. Radiology 1999, 213:341-6.
- el Mansari O, Sakit F, Janati Ml: Acute appendicitis on crural hernia. Presse Med 2002, 31(24):1129-30. French
- Guirguis EM, Taylor GA, Chadwick CD: Femoral appendicitis: an unusual case. Can J Surg 1989, 32(5):380-1.
- Naude GP, Ocon S, Bongard F: Femoral hernia: the dire consequences of a missed diagnosis. Am J Emerg Med 1997, 15(7):680-2.
- Wyatt JP, Varma JS: Femoral hernia appendix causing small intestinal obstruction. Postgrad Med J 1992, 68(797):223-4.
- Delamarre J, Descombes P, Grillot G, Deschepper B, Deramond H: Hydorcele of pancreatic origin. X-ray computed tomographic study of an intrascrotal collection in an acute outbreak of chronic pancreatitis. Radiol 1988, 69(11):689-90.
- Michalopoulos A, Papadopoulos V, Apostolitis S, Papavramidis T, Paramythiotis D, Berovalis P: A rare case of pancreatic pseudo-cyst masquerading as hydrocele. Acta Gastroenterol Belg 2006, 69:424.

Publish with **Bio Med Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours you keep the copyright

Submit your manuscript here: http://www.biomedcentral.com/info/publishing\_adv.asp

