

Case Report

Open Access

Small bowel obstruction secondary to migration of a fragment of lithobezoar: a case report

Mekki Medani*, Eddie Myers, Bryan Kenny and David Waldron

Address: Department of Surgery & Radiology, Mid-Western Regional Hospital, Limerick, Ireland

Email: Mekki Medani* - mekmed@gmail.com; Eddie Myers - mekmed@gmail.com; Bryan Kenny - mekmed@gmail.com; David Waldron - mekmed@gmail.com

* Corresponding author

Published: 7 December 2009

Received: 20 October 2009

Cases Journal 2009, 2:9155 doi:10.1186/1757-1626-2-9155

Accepted: 7 December 2009

This article is available from: <http://www.casesjournal.com/content/2/1/9155>

© 2009 Medani et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Introduction: Small bowel obstruction is a common world-wide condition that has a range of etiological factors. The management is largely dependent on the cause of the obstruction. Small bowel obstruction caused by foreign body ingestion is rare; many items have been reported as responsible, but there are no reports implicating polyurethane foam.

Case presentation: We report the case of a 44-year-old Irish male who presented following ingestion of polyurethane foam. He was asymptomatic on presentation but developed a small bowel obstruction shortly thereafter.

Conclusion: Patients presenting following ingestion of polyurethane foam should be scheduled for elective laparotomy, gastrotomy, and retrieval of the cast on the next available theatre list - given that they are suitable for surgery.

Case Presentation

We report the case of a 44-year-old gentleman with a background history of multiple laparotomies for deliberate self harm who presented six weeks following ingesting of polyurethane self-expanding foam. On admission, the patient had no signs or symptoms of small bowel obstruction. A plain film of his abdomen confirmed the presence of an intra-gastric cast of self-expanding foam (Figure 1).

A planned elective operation for laparotomy and retrieval of the cast was organized. In the interim, he went on to develop a small bowel obstruction. A CT scan of his abdomen confirmed the small bowel obstruction to be secondary to migration of a fragment of foam (Figure 2).

The obstruction settled with conservative management. Thereafter, he underwent a laparotomy, gastrotomy, and

removal of the cast on an elective basis (Figure 3). He made an uneventful post-operative recovery.

Discussion

Bowel obstruction was, until the late 19th and early 20th century, a concern of the physician rather than the surgeon. The value of surgical intervention was only recognized after the advent of anaesthesia and antisepsis.

In the developed world, intra-abdominal adhesions are recognized as the leading cause of small bowel obstruction, accounting for over 60% of all cases. Small bowel obstruction secondary to foreign bodies remains a rare cause.

Small bowel obstruction (SBO) secondary to foreign bodies (FB) is rare; it represents less than 6% of all SBO cases



Figure 1
Intra-gastric cast.

[1], and causes serious morbidity in less than one percent of all patients [2]. It is, however, commoner in certain cohorts, including children, and patients with alcoholism or psychosis [3]. Bezoar is the term used to describe the accumulation of undigested foreign bodies or food material in the gastrointestinal tract forming a conglomeration. The constituents of a bezoar generally dictate their




Figure 2
CT abdomen showing the large intra-gastric lithobezoar.



Figure 3
Cast following extraction.

nomenclature; e.g. phytobezoar (fibers or seeds of vegetables and fruits); trichobezoar (hair); lactobezoar (remnants of milk) and lithobezoar (rock or similar substances). Although they can be found in any part of the gastrointestinal system, the stomach is the most common site for bezoars [4].

Generally, bezoars remain in the stomach; however, they will occasionally pass to the small bowel [5]. Unlike in this case, initial radiographic imaging is frequently non-diagnostic with regards to the etiology of the obstruction, as the bezoars are not usually visible on plain films.

Other reported FB causing SBO include mesh plugs following open hernia repair, retained surgical gauze, magnetic toys, endoscopy capsules, retained PEG tube internal bumpers, gastric bands, and many others.

The signs and symptoms of FB SBO are the same as those of SBO secondary to any other cause, however a history of FB ingestion is usually evident; otherwise, a high index of suspicion is necessary, especially in children and patients with psychiatric illnesses.

In the management of FB ingestion, there are recommendations that are in place, recommended by a panel of experts [6]. These suggest that any foreign body that has not passed the stomach in three to four weeks should be removed endoscopically; which would not have been possible in this case. It also suggests that blunt objects beyond the stomach that remain in the same location for more than one week should be considered for surgical removal; and FB causing fever, vomiting, abdominal pain, or significant symptoms should be considered for emergency removal [6].

Conclusion

Patients presenting following ingestion of self-expanding foam should be scheduled for surgical retrieval on the next available elective list, as they are at risk of developing small bowel obstruction, due to the fragmentation of the lithobezoar.

Consent

Written informed consent was obtained from the patient for publication of this case report and accompanying images.

A copy of the written consent is available for review by the Editor-in-Chief of this journal.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

MM collected the patient information, performed the literature review, and was the major contributor in writing the manuscript. EM edited the manuscript and provided critical analysis of the body. BK interpreted the imaging and contributed to the image selection in the manuscript.

All authors have read and approved this final manuscript.

Acknowledgements

DW provided technical and writing assistance, and was of great support in finalizing the manuscript.

References

1. Richards WO, Williams LF Jr: **Obstruction of the large and small intestine.** *Surg Clin N Am* 1988:355-76.
2. Uyemura MC: **Foreign body ingestion in children.** *Am Fam Physician* 2005, **72(2)**:287-91.
3. Nijhawan S, Rai RR, Agarwal S, Vijayyergiya R: **Unusual foreign bodies of the upper gastrointestinal tract.** *Trop Gastroenterol* 1995:50-2.
4. Numanoglu KV, Tatli D: **A rare cause of intestinal obstruction in a child: colonic lithobezoar.** *Emerg Med J* 2008, **25(5)**:312-3.
5. Quintana JF, Walker RN, McGeehan A: **Child with small bowel obstruction and perforation secondary to ileal bezoar.** *Pediatr Emerg Care* 2008, **24(2)**:99-101.
6. Eisen GM, Baron TH, Dominitz JA, Faigel DO, Goldstein JL, Johanson JF, et al.: **Guideline for the management of ingested foreign bodies.** *Gastrointest Endosc* 2002:802-6.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

